



Murdoch, Craig and Kusch, J. and Flockhart, G. M. H. and Graham, D. and Faulds, K. and Uttamchandani, D. (2017) Effect of nanoparticle morphologies on signal strength in photoacoustic sensing. In: Optics + Ultrasound IV, 2017-11-23, University of Strathclyde. ,

This version is available at <https://strathprints.strath.ac.uk/64772/>

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: strathprints@strath.ac.uk

Effect of Nanoparticle Morphologies on Signal Strength in Photoacoustic Sensing

Craig Murdoch¹, J. Kusch¹, G. M. H. Flockhart¹, D. Graham², K. Faulds² and D. Uttamchandani¹

¹ *Department of Electrical and Electronic Engineering, University of Strathclyde*

² *Department of Pure & Applied Chemistry, University of Strathclyde*

Abstract

Photoacoustic sensing has the potential to probe to greater depths in tissues compared to optical microscopy. The use of plasmonic nanoparticles can further enhance the photoacoustic signal and the resonances of metallic nanoparticles, e.g. gold, can be aligned with commonly used visible and near-IR laser wavelengths. In addition, surface functionalised nanoparticles can be used to target specific biomolecules. In this work, we report the synthesis of spherical gold nanoparticles with a plasmonic extinction peak at 532 nm and two sizes of star shaped gold nanoparticles with extinction peaks at 532 nm and 600 nm. The nanoparticles were incorporated into tissue phantoms and the relative performance of the generation of photoacoustic signals from these different nanoparticle morphologies was investigated. At an excitation laser wavelength of 532 nm, we found that the spherical gold nanoparticles generated the greatest photoacoustic response.

Cite as: Murdoch, C., Kusch, J., Flockhart, G. M. H., Graham, D., Faulds, K., & Uttamchandani, D.(2017). *Effect of Nanoparticle Morphologies on Signal Strength in Photoacoustic Sensing*. Abstract from Optics + Ultrasound IV, Glasgow, United Kingdom.